

## POSTERIOR BOX COMPOSITE FINISHED RESTORATION

Ca	ADEX 2017  Tooth #: Examiner Number  Time Started:
	M D
	CRITICAL ERRORS
	The restoration is fractured
	ACC= Minimally Acceptable SUB= Marginally Substandard DEF= Critically Deficient
MAR	GIN INTEGRITY AND SURFACE FINISH
Margi	n Excess/Deficiency
ACC	There may be a marginal excess or deficiency at the restoration-tooth interface detectable either visually or with the tine of an explorer, but it is $\leq 0.5$ mm. There is no evidence of pits and/or voids at the cavosurface margin.
SUB	<ul> <li>A. DEFICIENCY: The restoration-tooth interface is detectable visually or with the tine of an explorer, and there is evidence of marginal deficiency &gt; 0.5 mm but ≤ 1.0 mm, which can include pits and/or voids at the cavosurface margin.</li> <li>B. EXCESS: The restoration-tooth interface is detectable visually or with the tine of an explorer, and there is evidence of marginal excess &gt; 0.5 mm but ≤ 1.0 mm, which can include pits and/or voids at the cavosurface margin. There is flash with contamination underneath, but it is not internal to the cavosurface margin and could be removed by polishing or finishing.</li> </ul>
DEF	<ul> <li>A. There is evidence of marginal deficiency of &gt; 1.0 mm, which includes pits and voids at the cavosurface margin, and/or there is an open margin.</li> <li>B. There is evidence of marginal excess (excluding bonding agent or unfilled resin) of &gt; 1.0 mm.</li> </ul>
Adjac	ent Tooth Structure
ACC	There is minimal or no evidence of unwarranted or unnecessary removal, modification, or recontouring of tooth structure adjacent to the restoration.
SUB	There is evidence of unwarranted or unnecessary removal, modification, or recontouring of tooth structure adjacent to the restoration.
DEF	There is gross enameloplasty resulting in the exposure of dentin.
Bondi	
ACC DEF	The restoration is bonded to the prepared tooth structure.  The restoration is debonded and/or movable in the preparation.
	TOUR, CONTACT AND OCCLUSION
Interp	proximal Contact
ACC	Interproximal contact is visually closed, and the contact is adequate in size, shape, or position but may demonstrate little resistance to dental floss.
DEF	A. The interproximal contact is visually open or concave/irregular, allowing for food impaction.  B. The interproximal contact will not allow floss to pass.

## Posterior Box Composite Finished Restoration - continued **Centric/Excursive Contacts** When checked with articulating paper, all centric and excursive contacts on the restoration are consistent in size, shape, and ACC intensity with such contacts on other teeth in that quadrant. When checked with articulating paper, the restoration is in hyper-occlusion, making it inconsistent in size, shape, and intensity **SUB** with the occlusal contacts on surrounding teeth, and it requires adjustment. There is gross hyper-occlusion so that the restoration is the only point of occlusion in that quadrant. TREATMENT MANAGEMENT **Adjacent Tooth Damage** Any damage to adjacent tooth/teeth can be removed with polishing without adversely altering the shape of the contour and/or ACC DEF There is evidence of gross damage and/or alteration to adjacent and/or opposing hard tissue inconsistent with the procedure. **Soft Tissue Damage ACC** The soft tissue is free from damage, or there is tissue damage that is consistent with the procedure. **DEF** There is gross iatrogenic damage to the soft tissue inconsistent with the procedure and pre-existing condition of the soft tissue.

Candidate Sequential		Examiner Number		
		Time Completed:		