

PREPARATION: ANTERIOR COMPOSITE

CDCA DENTAL THERAPY EXAM

Tooth #: _____

M	D	F	L
---	---	---	---

CRITICAL ERRORS

Wrong tooth/surface treated	No	Yes
Unrecognized exposure	No	Yes

ACC= Adheres to Criteria SUB= Marginally Substandard DEF= Critically Deficient

EXTERNAL OUTLINE FORM

Outline Extension

ACC	The wall opposite the access, if broken, may extend < 1.0 mm beyond the contact area. The outline form may be over-extended mesiodistally 0.5 mm to ≤ 1.0 mm beyond what is necessary for complete removal of caries and/or previous restorative material. The outline form dimension is ≤ 3.0 mm incisal gingivally.
SUB	A. The outline form is over-extended mesiodistally > 1.0 mm but ≤ 1.5 mm beyond what is necessary for complete removal of caries and/or previous restorative material. B. The incisal cavosurface margin is over-extended so that the integrity of the incisal angle is compromised. C. The wall opposite the access opening extends > 1.0 mm but ≤ 2.0 mm beyond the contact area. D. The outline form dimension is > 3.0 mm but ≤ 5.0 mm incisal gingivally.
DEF	A. The outline form is over-extended mesiodistally > 1.5 mm beyond necessary for complete removal of caries and/or previous restorative material. B. The incisal angle is unnecessarily removed or fractured. C. The wall opposite the access opening extends > 2.0 mm beyond the contact area. D. The outline form dimension is > 5.0 mm incisal gingivally. E. The outline form is under-extended, making it impossible to manipulate and finish the restorative material.

Gingival Contact Open

ACC	The gingival clearance is ≤ 1.0 mm.
SUB	A. The gingival clearance is > 1.0 mm but ≤ 2.0 mm.
DEF	A. The gingival clearance is > 2.0 mm.

Margin Smoothness/Continuity/Bevels

ACC	The cavosurface margins may be slightly irregular. Enamel cavosurface margin bevels, if present, are ≤ 1.0 mm in width.
SUB	The cavosurface margin is rough and severely irregular. Enamel cavosurface margin bevels, if present, are > 1.0 mm in width, are not uniform, or are inappropriate for the size of the restoration.

Sound Marginal Tooth Structure

ACC	There may be a small area of unsupported enamel which is not necessary to preserve facial aesthetics. There is no previous restorative material, excluding sealants, at the cavosurface margin.
SUB	A. There are large or multiple areas of unsupported enamel which are not necessary to preserve facial aesthetics. B. The cavosurface margin does not terminate in sound natural tooth structure. C. There is explorer-penetrable decalcification remaining on the cavosurface margin.

INTERNAL FORM**Axial Walls**

ACC	MAX ANTERIORS/MAND CUSPIDS: The depth of the axial wall extends ≤ 1.5 mm in depth from the DEJ. MAND INCISOR: The depth of the axial wall extends ≤ 1.25 mm in depth from the cavosurface margin.
SUB	MAX ANTERIORS/MAND CUSPIDS: The depth of the axial wall extends > 1.5 mm but ≤ 2.5 mm in depth from the DEJ. MAND INCISOR: The depth of the axial wall extends > 1.25 mm but ≤ 2.0 mm in depth from the cavosurface margin.
DEF	MAX ANTERIORS/MAND CUSPIDS: The depth of the axial wall extends > 2.5 mm in depth from the DEJ. MAND INCISOR: The depth of the axial wall extends > 2.0 mm beyond the cavosurface margin.

Internal Retention

ACC	If used, rounded internal retention is placed in the dentin of the gingival and incisal walls just axial to the DEJ as dictated by cavity form. Retention is tactilely and visually present.
SUB	When used, retention is excessive and undermines enamel, or jeopardizes the incisal angle, or encroaches on the pulp.

Caries/Remaining Material

ACC	All carious tooth structure and/or previous restorative material are removed.
DEF	A. Caries remain in the preparation. B. Previous restorative material remains in the preparation. C. Assigned carious lesions have not been accessed.

TREATMENT MANAGEMENT**Adjacent Tooth Damage**

ACC	Any damage to adjacent tooth/teeth can be removed with polishing without adversely altering the shape of the contour and/or contact.
DEF	There is gross damage to adjacent tooth/teeth which requires a restoration.

Soft Tissue Damage

ACC	The soft tissue is free from damage or there is tissue damage that is consistent with the procedure.
DEF	There is gross iatrogenic damage to the soft tissue inconsistent with the procedure and pre-existing condition of the soft tissue.